

8月31日 までは本年値、それ以降は平年値を使用

		収 穫 期																																																													
		8/15	8/16	8/17	8/18	8/19	8/20	8/21	8/22	8/23	8/24	8/25	8/26	8/27	8/28	8/29	8/30	8/31	9/1	9/2	9/3	9/4	9/5	9/6	9/7	9/8	9/9	9/10	9/11	9/12	9/13	9/14	9/15	9/16	9/17	9/18	9/19	9/20	9/21	9/22	9/23	9/24	9/25																				
平均 気温	本年値	27.5	27.6	28.0	28.1	28.5	27.9	25.9	26.5	27.6	29.0	29.9	28.1	22.9	23.2	20.5	25.6	27.1																																													
	平年値	26.6	26.5	26.5	26.4	26.3	26.2	26.1	26.0	25.9	25.8	25.7	25.6	25.5	25.4	25.3	25.1	25.0	24.9	24.8	24.7	24.5	24.4	24.2	24.0	23.8	23.6	23.5	23.3	23.1	22.9	22.7	22.5	22.3	22.0	21.8	21.6	21.4	21.2	21.0	20.8	20.6	20.4																				
出 穂 期	7/10	992	1020	1048	1076	1104	1132	1158	1185	1,212	1,241																			出穂期と収穫適期のめやす(平坦地)																																	
	7/11	965	993	1021	1049	1077	1105	1131	1158	1185	1,214	1,244																																																			
	7/12	938	965	993	1022	1050	1078	1104	1130	1158	1187	1,217	1,245																																																		
	7/13	909	936	964	992	1021	1049	1075	1101	1129	1158	1188	1,216	1,239																																																	
	7/14	881	908	936	964	993	1021	1047	1073	1101	1130	1160	1188	1,211	1,234																																																
	7/15	852	879	907	935	964	992	1018	1044	1072	1101	1131	1159	1182	1,205	1,225																																															
	7/16	824	852	880	908	936	964	990	1017	1044	1073	1103	1131	1154	1,177	1,198	1,223																																														
	7/17	797	825	853	881	909	937	963	990	1017	1046	1076	1104	1127	1,150	1,171	1,196	1,223	1,248																																												
	7/18		797	825	853	882	909	935	962	989	1018	1048	1076	1099	1,123	1,143	1,169	1,196	1,221	1,245																																											
	7/19			796	824	852	880	906	933	960	989	1019	1047	1070	1,093	1,114	1,139	1,167	1,191	1,216	1,241																																										
	7/20				794	822	850	876	903	930	959	989	1017	1040	1,063	1,084	1,109	1,137	1,161	1,186	1,211	1,235																																									
	7/21				765	793	821	847	874	901	930	960	988	1,011	1,034	1,055	1,080	1,108	1,132	1,157	1,182	1,206	1,231																																								
	7/22					792	818	844	872	901	931	959	982	1,005	1,025	1,051	1,078	1,103	1,128	1,152	1,177	1,201	1,225	1,249																																							
	7/23						789	815	843	872	902	930	953	976	996	1,022	1,049	1,074	1,099	1,124	1,148	1,172	1,197	1,221	1,244																																						
	7/24								814	843	873	901	924	947	968	993	1,020	1,045	1,070	1,095	1,119	1,144	1,168	1,192	1,216	1,239																																					
	7/25									816	846	874	897	920	940	966	993	1,018	1,043	1,067	1,092	1,116	1,140	1,164	1,188	1,212	1,235																																				
	7/26										821	849	872	895	916	942	969	994	1,018	1,043	1,068	1,092	1,116	1,140	1,164	1,188	1,211	1,234																																			
	7/27											824	847	870	891	916	943	968	993	1,018	1,042	1,067	1,091	1,115	1,139	1,162	1,186	1,209	1,232																																		
	7/28												822	845	866	891	918	943	968	993	1,017	1,042	1,066	1,090	1,114	1,137	1,161	1,184	1,207	1,230																																	
	7/29													820	840	866	893	918	943	967	992	1,016	1,040	1,064	1,088	1,112	1,135	1,159	1,182	1,205	1,227	1,250																															
7/30														813	838	865	890	915	940	964	989	1,013	1,037	1,061	1,084	1,108	1,131	1,154	1,177	1,200	1,222	1,245																															
7/31															810	837	862	887	912	936	961	985	1,009	1,033	1,056	1,080	1,103	1,126	1,149	1,172	1,194	1,217	1,239																														
8/1																	810	834	859	884	908	933	957	981	1,005	1,028	1,052	1,075	1,098	1,121	1,144	1,166	1,189	1,211	1,233																												
8/2																		807	832	857	881	906	930	954	978	1,001	1,025	1,048	1,071	1,094	1,117	1,139	1,162	1,184	1,205	1,227	1,248																										
8/3	収穫適期のめやす（出穂後の積算温度・通常年の場合） 【早生品種】 なつひめ 1, 000℃前後 五百万石、こしいぶき 975℃ つきあかり 1, 000～1, 100℃ 【中生品種】 こがねもち、コシヒカリ . . . 1, 000℃ 【晩生品種】 新之助 1, 050～1, 100℃ ○高温登熟条件となる場合には、積算温度を上記のめやすより50℃（2日程度） 早めましょう。 ○実際の収穫は粳の黄化割合で判断しましょう。																			804	829	854	878	902	926	950	974	997	1,020	1,043	1,066	1,089	1,112	1,134	1,156	1,178	1,199	1,221	1,242																								
8/4																																				801	826	850	874	898	922	946	969	992	1,016	1,038	1,061	1,084	1,106	1,128	1,150	1,171	1,193	1,214	1,235								
8/5																																						822	846	870	894	917	941	964	987	1,010	1,033	1,055	1,078	1,100	1,122	1,143	1,165	1,186	1,207	1,228	1,248						
8/6																																								817	841	865	889	912	935	958	981	1,004	1,027	1,049	1,071	1,093	1,114	1,136	1,157	1,178	1,199	1,219	1,240				
8/7																																										811	835	859	882	906	929	952	974	997	1,019	1,041	1,063	1,085	1,106	1,127	1,148	1,169	1,190	1,210			
8/8																																												807	831	854	877	901	923	946	969	991	1,013	1,035	1,056	1,078	1,099	1,120	1,141	1,161	1,182		
8/9																																													804	827	851	874	897	919	942	964	986	1,008	1,029	1,051	1,072	1,093	1,114	1,134	1,155		
8/10																																														802	825	848	871	894	916	938	960	982	1,004	1,025	1,046	1,067	1,088	1,109	1,129		
8/11																																															801	824	847	870	893	915	937	959	980	1,002	1,023	1,044	1,065	1,085	1,106		
8/12																																																															
8/13																																																															
8/14																																																															
8/15																																																															